


REPORT DOCUMENTATION PAGE				<i>Form Approved</i> OMB No. 0704-0188	
Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number. PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.					
1. REPORT DATE (DD-MM-YYYY) 21-08-2003		2. REPORT TYPE Technical Viewgraph Presentation		3. DATES COVERED (From - To)	
4. TITLE AND SUBTITLE New Ionic Liquids				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S) Greg Drake, Tom Hawkins (AFRL/PRSP); Kerri Tollison (ERC); Leslie Hall, Ashwani Vij (AFRL/PRSP)				5d. PROJECT NUMBER 2303	
				5e. TASK NUMBER M2C8	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Air Force Research Laboratory (AFMC) AFRL/PRS 5 Pollux Drive Edwards AFB CA 93524-7048				8. PERFORMING ORGANIZATION REPORT NUMBER AFRL-PR-ED-VG-2003-210	
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) Air Force Research Laboratory (AFMC) AFRL/PRS 5 Pollux Drive Edwards AFB CA 93524-7048				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S NUMBER(S) AFRL-PR-ED-VG-2003-210	
12. DISTRIBUTION / AVAILABILITY STATEMENT Approved for public release; distribution unlimited.					
13. SUPPLEMENTARY NOTES For presentation at the National ACS Meeting (Ionic Liquids Symposium) in New York, NY, taking place 7-11 September 2003.					
14. ABSTRACT					
20030929 087					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT	18. NUMBER OF PAGES	19a. NAME OF RESPONSIBLE PERSON
a. REPORT Unclassified	b. ABSTRACT Unclassified	c. THIS PAGE Unclassified	A	27	Leilani Richardson
					19b. TELEPHONE NUMBER (include area code) (661) 275-5015

New Ionic Liquids

**ACS Ionic Liquids Symposium
September 9, 2003**




**Greg Drake, Tommy Hawkins, Kerri Tollison*,
Leslie Hall, Ashwani Vij, Sarah Sabowski*
AFRL/PRSP and *ERC, Inc.
Air Force Research Laboratory
Edwards Air Force Base, CA 93524-7680**

Distribution Statement A: Public Release, Distribution unlimited.



New Ionic Liquids



What are Ionic Liquids?

A class of salts consisting of cation/anion pair that has a very low melting point.


(1) An ionic compound that melts below 100 °C (b.p. of H₂O). J. Wilkes, P. Wasserscheid, K. Seddon.

(2) An ionic compound that has a melting point at or below ambient temperatures. These are often called RTILs (Room Temperature Ionic Liquids) T. Welton, R. Rogers.

But many of the salts fit both definitions and 2 is really a more specific class of (1), and all are low melting salts.



New Ionic Liquids




Important factors affecting the physical properties of ionic liquids


1. Asymmetry of cation as well as anion
2. Packing efficiency
3. Charge delocalization in cationic/anionic species
4. “Sheer size” differentials



New Ionic Liquids




Ionic liquids research was really opened up by the pioneering work of King, Wilkes, and Hussey under USAF research looking for new battery electrolytes.¹ Dealt heavily with aluminum halide anions early on.



1-ethyl-3-methylimidazolium tetrachloroaluminate

Later, water stable ionic liquids were synthesized by Wilkes and coworkers which opened up the currently rapidly expanding field that we see today².




1-butyl-3-methylimidazolium hexafluorophosphate

¹Fannin, A. ; Floreani, D. ; King, L. ; Landers, J. ; Piersma, B. ; Stech, D. ; Vaughn, R. ; Wilkes, J.S. *J. Phys. Chem.* **1984**, *88*, 2614. ²Wilkes, J. *J. Chem. Soc. Chem. Commun.* **1992**, 965.




New Ionic Liquids




Wasserscheid, P.; Keim, W. *Angew. Chem. Int. Ed. Engl.* 2000, 39, 3772. Wasserscheid, P., Welton, T. (eds.) *Ionic Liquids in Synthesis* Wiley-VCH, FRG, 2003. Seddon, K.R.; Holbrey, J.D. *Clean Products and Processes* 1999, 1, 223. Rogers, R.; Seddon, K. (eds.) *Ionic Liquids* A.C.S. Symp. Ser. 818 2002 A.C.S. Publ. Co.




New Ionic Liquids



Most ionic liquids are based upon imidazolium rings and “heavy” or “dead” anions. We felt that we could use the shape of the cation and the poor fit idea to make much more energetic salts in a simple manner.



1-n-butyl-3-methyl imidazolium
cation




1-ethyl-4-amino-1,2,4-
triazolium cation

These new ionic liquids have similar shapes and physical properties, BUT higher ΔH_f , higher densities, and better oxygen balances.




New Ionic Liquids




4-amino-1,2,4-triazole

+ XS



N-propyl bromide




**Synthesis is from commercial materials
High yield simple isolation has been known
in literature for quite sometime.**

1-n-propyl-4-amino-
1,2,4-triazolium bromide
(yield >95% very pure)


Scriven; Keay; Goe; Astleford J. Org. Chem. **1989**, *54*, 731.




New Ionic Liquids




1-n-butyl-4-amino-1,2,4-triazolium cation




1-(2-ethanol)-4-amino-1,2,4-triazolium cation




1-isopropyl-4-amino-1,2,4-triazolium cation





1-methylcyclopropyl-4-amino-1,2,4-triazolium cation




1-(2-aminoethyl)-4-amino-1,2,4-triazolium dication




New Ionic Liquids




1-methyl-4-amino-1,2,4-triazolium cation



1-ethyl-4-amino-1,2,4-triazolium cation




1-(2-propenyl)-4-amino-1,2,4-triazolium cation



1-n-propyl-4-amino-1,2,4-triazolium cation



New Ionic Liquids



Physical properties of 1-n-alkyl substituted-4-amino-1,2,4-triazolium bromides.

- increasing melting points with increasing molecular weights,
- decomposition onsets that are relatively low
- densities decrease with increasing alkyl chain length.


Substituted 4AT salts	m.p. (°C)	dec. onset (°C)	density (g/cm ³)
1-ethyl	63°	110	1.69
1-n-propyl	60°	120	1.56
1-isopropyl	90°	110	1.60
1-butyl	48°	130	1.46
1-n-pentyl	54°	130	1.37
1-n-hexyl	76°	120	1.34
1-n-heptyl	94°	120	1.30
1-n-octyl	80°	135	1.27
1-n-nonyl	81°	140	1.26
1-n-decyl	90°	135	1.23



New Ionic Liquids




proton of 1-n-butyl-4-amino-1,2,4-triazolium bromide solid in d₆-DMSO



Current Data Parameters
NAME 4m-butyl-Salt
EXPNO 1
PROCNO 1
F2-Acquisition Parameters
Date_ 20020725
Time 10.46
INSTRUM spect
PROBHD 6mm BBO BP-
PULPROG zgpg
TD 65536
SOLVENT DMSO
NS 32
DS 2
SWH 8278.146 Hz
FIDRES 0.126314 Hz
AQ 3.869440 sec
RG 32
DW 60.00 usec
DE 6.00 usec
TE 300.2 K
DT 1.0000000 sec
===== CHANNEL f1 =====
NUC1 1H
P1 7.20 usec
PL1 -8.00 dB
SFO1 400.124710 MHz
F2-Processing parameters
SI 32768
SF 400.1300000 MHz
WDW EM
SSB 0
LB 0.30 Hz
GB 0
PC 1.00


carbon of 1-n-butyl-4-amino-1,2,4-triazolium bromide solid in d₆-DMSO



¹H(left) and ¹³C nmr spectra of 1-butyl-4-amino-1,2,4-triazolium bromide.





New Ionic Liquids



Single x-ray diffraction study of 1-ethyl-4-amino-1,2,4-triazolium bromide.



New Ionic Liquids





Single crystal x-ray diffraction study of 1-n-propyl-4-amino-1,2,4-triazolium bromide showing significant hydrogen bond contacts.

[illegible]

47




New Ionic Liquids



Single crystal x-ray diffraction structure of 1-isopropyl-4-amino-1,2,4-triazolium bromide.




New Ionic Liquids




Single crystal x-ray diffraction study of 1-hexyl-4-amino-1,2,4-triazolium bromide.




New Ionic Liquids



Single crystal x-ray diffraction study of 1-heptyl-4-amino-1,2,4-triazolium bromide.





New Ionic Liquids



Hydrogen bond contacts in 1-heptyl-4-amino-1,2,4-triazolium bromide



New Ionic Liquids



1-dodecyl-3-methylimidazolium hexafluorophosphate*

1-hexyl-4-amino-1,2,4-triazolium bromide#

*Gordon, C. M.; Holbrey, J. D.; Kennedy, A. R.; Seddon, K. R. *J. Mater. Chem.* **1998**, *8*, 2627. #Drake, G. W.; Hawkins, T. W.; Tollison, K.; Hall, L.; Vij, A. 2003 manuscript in progress.




New Ionic Liquids



But halides are only the start...

Nitrates were best made through silver nitrate metathesis in methanol.



This route led to the best materials as the silver bromide was easily removed.



New Ionic Liquids



1-substituted-4-amino-1,2,4-triazolium nitrate salts are more stable.

<u>Salt</u>	<u>melting point(°C)</u>	<u>decomp onset(°C)</u>	<u>$\rho(\text{g}/\text{cm}^3, \text{est.})$</u>
1-methyl	54	185	1.57
1-ethyl	5	185	1.39 (1.38)
1-n-propyl	34	190	1.35
1-isopropyl	53	175	1.37 (1.43)
1-n-butyl	-10	190	1.31
1-(2-ethanol)	10	180	1.48
1-methylcyclopropyl	56	190	1.36 (1.44)
1-(2-propenyl)	10	165	1.23
1-n-pentyl	26	170	1.29
1-n-hexyl	-2	160	1.26
1-n-heptyl	31	160	1.24
1-n-octyl	29	170	1.22
1-n-nonyl	53	175	1.20
1-n-decyl	49	185	1.18




New Ionic Liquids



Sample: 1-PROPYL-4-AT NITRATE
Size: 1.9000 mg
Method: greg
Comment: 10C/min/10ml/minhermeticalpans

DSC



File: C:\...files from old DSC\4at propyl no3
Operator: DRAKE
Run Date: 16-Jan-02 23:04



DSC of 1-n-propyl-4-amino-1,2,4-triazolium nitrate




New Ionic Liquids



Single crystal x-ray diffraction study of 1-methylcyclopropyl-4-amino-1,2,4-triazolium nitrate.



New Ionic Liquids




Single crystal x-ray diffraction structure of 1-isopropyl-4-amino-1,2,4-triazolium nitrate



New Ionic Liquids



The new energetic cations are weakly acidic in nature, aqueous solutions have a pH of around 4 which suggests the equilibrium involving a zwitterionic 1-alkyl-4-amido-1,2,4-triazolium species. This equilibrium could be one possible way for the ionic liquids to “come apart”.





New Ionic Liquids



Summary and Conclusions

A large new class of low melting salts which should be considered as new members of the well known class of materials referred to as ionic liquids has been synthesized and well characterized.

Using asymmetric cation shapes and poor cation-anion fit, an analogue system to the well known 1,3-dialkylsubstituted imidazolium cation family, based upon 1-substituted-4-amino-1,2,4-triazolium cations paired with the bromide and nitrate ions has been explored.

Facile synthesis routes from commercially available materials coupled with high yield and purity reactions make these new materials very exciting.


Several single crystal x-ray diffraction studies of several structures have been carried out proving the expected structure as well as revealing extensive hydrogen bonding in the solid state.

Physical properties of 1-substituted-4-amino-1,2,4-triazolium salts included much higher viscosities, higher densities, and much more polar behavior than that of imidazolium ionic liquids.

Further work is being carried out with other ions.



New Ionic Liquids



ACKNOWLEDGEMENTS

- MIKE BERMAN (AFOSR)
- WAYNE KALLIOMAA; RONALD CHANNELL (AFRL/PRSP)
- JOHN WILKES (USAF)
- JEFF SHEEHY (NASA/MARSHALL AND UA/HUNTSVILLE)
- CLAUDE MERRILL
- TOMMY HIGHSMITH
- JEFF BOTTARO, MARK PETRIE (SRI, INT.)
- MIKE HUGGINS (AFRL SUPPORT)